

8.3 Trigonometric Integrals

- Solve trigonometric integrals involving powers of sine and cosine.
- Solve trigonometric integrals involving powers of secant and tangent.
- Solve trigonometric integrals involving sine-cosine products with different angles.

Integrals Involving Powers of Sine and Cosine

In this section you will study techniques for evaluating integrals of the form

$$\int \sin^m x \cos^n x \, dx \quad \text{and} \quad \int \sec^m x \tan^n x \, dx$$

where either m or n is a positive integer. To find antiderivatives for these forms, try to break them into combinations of trigonometric integrals to which you can apply the Power Rule.

For instance, you can evaluate $\int \sin^5 x \cos x \, dx$ with the Power Rule by letting $u = \sin x$. Then, $du = \cos x \, dx$ and you have

$$\int \sin^5 x \cos x \, dx = \int u^5 \, du = \frac{u^6}{6} + C = \frac{\sin^6 x}{6} + C.$$

To break up $\int \sin^m x \cos^n x \, dx$ into forms to which you can apply the Power Rule, use the following identities.

$$\begin{aligned} \sin^2 x + \cos^2 x &= 1 && \text{Pythagorean identity} \\ \sin^2 x &= \frac{1 - \cos 2x}{2} && \text{Half-angle identity for } \sin^2 x \\ \cos^2 x &= \frac{1 + \cos 2x}{2} && \text{Half-angle identity for } \cos^2 x \end{aligned}$$

SHEILA SCOTT MACINTYRE (1910–1960)

Sheila Scott Macintyre published her first paper on the asymptotic periods of integral functions in 1935. She completed her doctorate work at Aberdeen University, where she taught. In 1958 she accepted a visiting research fellowship at the University of Cincinnati.

GUIDELINES FOR EVALUATING INTEGRALS INVOLVING POWERS OF SINE AND COSINE

1. If the power of the sine is odd and positive, save one sine factor and convert the remaining factors to cosines. Then, expand and integrate.

$$\int \overbrace{\sin^{2k+1} x}^{\text{Odd}} \cos^n x \, dx = \int \overbrace{(\sin^2 x)^k}^{\text{Convert to cosines}} \overbrace{\cos^n x \sin x}^{\text{Save for } du} \, dx = \int (1 - \cos^2 x)^k \cos^n x \sin x \, dx$$

2. If the power of the cosine is odd and positive, save one cosine factor and convert the remaining factors to sines. Then, expand and integrate.

$$\int \sin^m x \overbrace{\cos^{2k+1} x}^{\text{Odd}} \, dx = \int \sin^m x \overbrace{(\cos^2 x)^k}^{\text{Convert to sines}} \overbrace{\cos x}^{\text{Save for } du} \, dx = \int \sin^m x (1 - \sin^2 x)^k \cos x \, dx$$

3. If the powers of both the sine and cosine are even and nonnegative, make repeated use of the identities

$$\sin^2 x = \frac{1 - \cos 2x}{2} \quad \text{and} \quad \cos^2 x = \frac{1 + \cos 2x}{2}$$

to convert the integrand to odd powers of the cosine. Then proceed as in guideline 2.

TECHNOLOGY Use a computer algebra system to find the integral in Example 1. You should obtain

$$\int \sin^3 x \cos^4 x \, dx = -\cos^5 x \left(\frac{1}{7} \sin^2 x + \frac{2}{35} \right) + C.$$

Is this equivalent to the result obtained in Example 1?

EXAMPLE 1 Power of Sine Is Odd and Positive

Find $\int \sin^3 x \cos^4 x \, dx$.

Solution Because you expect to use the Power Rule with $u = \cos x$, save one sine factor to form du and convert the remaining sine factors to cosines.

$$\begin{aligned} \int \sin^3 x \cos^4 x \, dx &= \int \sin^2 x \cos^4 x (\sin x) \, dx && \text{Rewrite.} \\ &= \int (1 - \cos^2 x) \cos^4 x \sin x \, dx && \text{Trigonometric identity} \\ &= \int (\cos^4 x - \cos^6 x) \sin x \, dx && \text{Multiply.} \\ &= \int \cos^4 x \sin x \, dx - \int \cos^6 x \sin x \, dx && \text{Rewrite.} \\ &= -\int \cos^4 x (-\sin x) \, dx + \int \cos^6 x (-\sin x) \, dx \\ &= -\frac{\cos^5 x}{5} + \frac{\cos^7 x}{7} + C && \text{Integrate.} \end{aligned}$$


In Example 1, both of the powers m and n happened to be positive integers. However, the same strategy will work as long as either m or n is odd and positive. For instance, in the next example the power of the cosine is 3, but the power of the sine is $-\frac{1}{2}$.

EXAMPLE 2 Power of Cosine Is Odd and Positive

Evaluate $\int_{\pi/6}^{\pi/3} \frac{\cos^3 x}{\sqrt{\sin x}} \, dx$.

Solution Because you expect to use the Power Rule with $u = \sin x$, save one cosine factor to form du and convert the remaining cosine factors to sines.

$$\begin{aligned} \int_{\pi/6}^{\pi/3} \frac{\cos^3 x}{\sqrt{\sin x}} \, dx &= \int_{\pi/6}^{\pi/3} \frac{\cos^2 x \cos x}{\sqrt{\sin x}} \, dx \\ &= \int_{\pi/6}^{\pi/3} \frac{(1 - \sin^2 x)(\cos x)}{\sqrt{\sin x}} \, dx \\ &= \int_{\pi/6}^{\pi/3} [(\sin x)^{-1/2} \cos x - (\sin x)^{3/2} \cos x] \, dx \\ &= \left[\frac{(\sin x)^{1/2}}{1/2} - \frac{(\sin x)^{5/2}}{5/2} \right]_{\pi/6}^{\pi/3} \\ &= 2 \left(\frac{\sqrt{3}}{2} \right)^{1/2} - \frac{2}{5} \left(\frac{\sqrt{3}}{2} \right)^{5/2} - \sqrt{2} + \frac{\sqrt{32}}{80} \\ &\approx 0.239 \end{aligned}$$



The area of the region is approximately 0.239.

Figure 8.4

Figure 8.4 shows the region whose area is represented by this integral.

EXAMPLE 3 Power of Cosine Is Even and Nonnegative

Find $\int \cos^4 x \, dx$.

Solution Because m and n are both even and nonnegative ($m = 0$), you can replace $\cos^4 x$ by $[(1 + \cos 2x)/2]^2$.


$$\begin{aligned} \int \cos^4 x \, dx &= \int \left(\frac{1 + \cos 2x}{2} \right)^2 dx && \text{Half-angle identity} \\ &= \int \left(\frac{1}{4} + \frac{\cos 2x}{2} + \frac{\cos^2 2x}{4} \right) dx && \text{Expand.} \\ &= \int \left[\frac{1}{4} + \frac{\cos 2x}{2} + \frac{1}{4} \left(\frac{1 + \cos 4x}{2} \right) \right] dx && \text{Half-angle identity} \\ &= \frac{3}{8} \int dx + \frac{1}{4} \int 2 \cos 2x \, dx + \frac{1}{32} \int 4 \cos 4x \, dx && \text{Rewrite.} \\ &= \frac{3x}{8} + \frac{\sin 2x}{4} + \frac{\sin 4x}{32} + C && \text{Integrate.} \end{aligned}$$

Use a symbolic differentiation utility to verify this. Can you simplify the derivative to obtain the original integrand? ■

In Example 3, if you were to evaluate the definite integral from 0 to $\pi/2$, you would obtain

$$\begin{aligned} \int_0^{\pi/2} \cos^4 x \, dx &= \left[\frac{3x}{8} + \frac{\sin 2x}{4} + \frac{\sin 4x}{32} \right]_0^{\pi/2} \\ &= \left(\frac{3\pi}{16} + 0 + 0 \right) - (0 + 0 + 0) \\ &= \frac{3\pi}{16} \end{aligned}$$

Note that the only term that contributes to the solution is $3x/8$. This observation is generalized in the following formulas developed by John Wallis.



Bettmann/Corbis

JOHN WALLIS (1616–1703)

Wallis did much of his work in calculus prior to Newton and Leibniz, and he influenced the thinking of both of these men. Wallis is also credited with introducing the present symbol (∞) for infinity.

WALLIS'S FORMULAS

1. If n is odd ($n \geq 3$), then

$$\int_0^{\pi/2} \cos^n x \, dx = \left(\frac{2}{3} \right) \left(\frac{4}{5} \right) \left(\frac{6}{7} \right) \cdots \left(\frac{n-1}{n} \right).$$
2. If n is even ($n \geq 2$), then

$$\int_0^{\pi/2} \cos^n x \, dx = \left(\frac{1}{2} \right) \left(\frac{3}{4} \right) \left(\frac{5}{6} \right) \cdots \left(\frac{n-1}{n} \right) \left(\frac{\pi}{2} \right).$$

These formulas are also valid if $\cos^n x$ is replaced by $\sin^n x$. (You are asked to prove both formulas in Exercise 108.)

Integrals Involving Powers of Secant and Tangent

The following guidelines can help you evaluate integrals of the form

$$\int \sec^m x \tan^n x \, dx.$$

GUIDELINES FOR EVALUATING INTEGRALS INVOLVING POWERS OF SECANT AND TANGENT

1. If the power of the secant is even and positive, save a secant-squared factor and convert the remaining factors to tangents. Then expand and integrate.

$$\int \overbrace{\sec^{2k} x}^{\text{Even}} \tan^n x \, dx = \int \overbrace{(\sec^2 x)^{k-1}}^{\text{Convert to tangents}} \overbrace{\tan^n x \sec^2 x}^{\text{Save for } du} \, dx = \int (1 + \tan^2 x)^{k-1} \tan^n x \sec^2 x \, dx$$

2. If the power of the tangent is odd and positive, save a secant-tangent factor and convert the remaining factors to secants. Then expand and integrate.

$$\int \sec^m x \overbrace{\tan^{2k+1} x}^{\text{Odd}} \, dx = \int \sec^{m-1} x \overbrace{(\tan^2 x)^k}^{\text{Convert to secants}} \overbrace{\sec x \tan x}^{\text{Save for } du} \, dx = \int \sec^{m-1} x (\sec^2 x - 1)^k \sec x \tan x \, dx$$

3. If there are no secant factors and the power of the tangent is even and positive, convert a tangent-squared factor to a secant-squared factor, then expand and repeat if necessary.

$$\int \tan^n x \, dx = \int \tan^{n-2} x \overbrace{(\tan^2 x)}^{\text{Convert to secants}} \, dx = \int \tan^{n-2} x (\sec^2 x - 1) \, dx$$

4. If the integral is of the form $\int \sec^m x \, dx$, where m is odd and positive, use integration by parts, as illustrated in Example 5 in the preceding section.
5. If none of the first four guidelines applies, try converting to sines and cosines.

EXAMPLE 4 Power of Tangent Is Odd and Positive

Find $\int \frac{\tan^3 x}{\sqrt{\sec x}} \, dx$.

Solution Because you expect to use the Power Rule with $u = \sec x$, save a factor of $(\sec x \tan x)$ to form du and convert the remaining tangent factors to secants.

$$\begin{aligned} \int \frac{\tan^3 x}{\sqrt{\sec x}} \, dx &= \int (\sec x)^{-1/2} \tan^3 x \, dx \\ &= \int (\sec x)^{-3/2} (\tan^2 x) (\sec x \tan x) \, dx \\ &= \int (\sec x)^{-3/2} (\sec^2 x - 1) (\sec x \tan x) \, dx \\ &= \int [(\sec x)^{1/2} - (\sec x)^{-3/2}] (\sec x \tan x) \, dx \\ &= \frac{2}{3} (\sec x)^{3/2} + 2(\sec x)^{-1/2} + C \end{aligned}$$

NOTE In Example 5, the power of the tangent is odd and positive. So, you could also find the integral using the procedure described in guideline 2 on page 539. In Exercise 89, you are asked to show that the results obtained by these two procedures differ only by a constant.

EXAMPLE 5 Power of Secant Is Even and Positive

Find $\int \sec^4 3x \tan^3 3x \, dx$.

Solution Let $u = \tan 3x$, then $du = 3 \sec^2 3x \, dx$ and you can write

$$\begin{aligned} \int \sec^4 3x \tan^3 3x \, dx &= \int \sec^2 3x \tan^3 3x (\sec^2 3x) \, dx \\ &= \int (1 + \tan^2 3x) \tan^3 3x (\sec^2 3x) \, dx \\ &= \frac{1}{3} \int (\tan^3 3x + \tan^5 3x) (3 \sec^2 3x) \, dx \\ &= \frac{1}{3} \left(\frac{\tan^4 3x}{4} + \frac{\tan^6 3x}{6} \right) + C \\ &= \frac{\tan^4 3x}{12} + \frac{\tan^6 3x}{18} + C. \end{aligned}$$

EXAMPLE 6 Power of Tangent Is Even

Evaluate $\int_0^{\pi/4} \tan^4 x \, dx$.


Solution Because there are no secant factors, you can begin by converting a tangent-squared factor to a secant-squared factor.

$$\begin{aligned} \int \tan^4 x \, dx &= \int \tan^2 x (\tan^2 x) \, dx \\ &= \int \tan^2 x (\sec^2 x - 1) \, dx \\ &= \int \tan^2 x \sec^2 x \, dx - \int \tan^2 x \, dx \\ &= \int \tan^2 x \sec^2 x \, dx - \int (\sec^2 x - 1) \, dx \\ &= \frac{\tan^3 x}{3} - \tan x + x + C \end{aligned}$$

You can evaluate the definite integral as follows.

$$\begin{aligned} \int_0^{\pi/4} \tan^4 x \, dx &= \left[\frac{\tan^3 x}{3} - \tan x + x \right]_0^{\pi/4} \\ &= \frac{\pi}{4} - \frac{2}{3} \\ &\approx 0.119 \end{aligned}$$

The area represented by the definite integral is shown in Figure 8.5. Try using Simpson's Rule to approximate this integral. With $n = 18$, you should obtain an approximation that is within 0.00001 of the actual value. ■



The area of the region is approximately 0.119.

Figure 8.5

For integrals involving powers of cotangents and cosecants, you can follow a strategy similar to that used for powers of tangents and secants. Also, when integrating trigonometric functions, remember that it sometimes helps to convert the entire integrand to powers of sines and cosines.

EXAMPLE 7 Converting to Sines and Cosines

Find $\int \frac{\sec x}{\tan^2 x} dx$.

Solution Because the first four guidelines on page 539 do not apply, try converting the integrand to sines and cosines. In this case, you are able to integrate the resulting powers of sine and cosine as follows.

$$\begin{aligned}\int \frac{\sec x}{\tan^2 x} dx &= \int \left(\frac{1}{\cos x} \right) \left(\frac{\cos x}{\sin x} \right)^2 dx \\ &= \int (\sin x)^{-2} (\cos x) dx \\ &= -(\sin x)^{-1} + C \\ &= -\csc x + C\end{aligned}$$

Integrals Involving Sine-Cosine Products with Different Angles

FOR FURTHER INFORMATION

To learn more about integrals involving sine-cosine products with different angles, see the article “Integrals of Products of Sine and Cosine with Different Arguments” by Sherrie J. Nicol in *The College Mathematics Journal*. To view this article, go to the website www.matharticles.com.

Integrals involving the products of sines and cosines of two *different* angles occur in many applications. In such instances you can use the following product-to-sum identities.

$$\sin mx \sin nx = \frac{1}{2}(\cos[(m-n)x] - \cos[(m+n)x])$$

$$\sin mx \cos nx = \frac{1}{2}(\sin[(m-n)x] + \sin[(m+n)x])$$

$$\cos mx \cos nx = \frac{1}{2}(\cos[(m-n)x] + \cos[(m+n)x])$$

EXAMPLE 8 Using Product-to-Sum Identities

Find $\int \sin 5x \cos 4x dx$.

Solution Considering the second product-to-sum identity above, you can write

$$\begin{aligned}\int \sin 5x \cos 4x dx &= \frac{1}{2} \int (\sin x + \sin 9x) dx \\ &= \frac{1}{2} \left(-\cos x - \frac{\cos 9x}{9} \right) + C \\ &= -\frac{\cos x}{2} - \frac{\cos 9x}{18} + C.\end{aligned}$$

8.3 Exercises

See www.CalcChat.com for worked-out solutions to odd-numbered exercises.

In Exercises 1–4, use differentiation to match the antiderivative with the correct integral. [Integrals are labeled (a), (b), (c), and (d).]

(a) $\int \sin x \tan^2 x \, dx$ (b) $8 \int \cos^4 x \, dx$

(c) $\int \sin x \sec^2 x \, dx$ (d) $\int \tan^4 x \, dx$

1. $y = \sec x$
2. $y = \cos x + \sec x$
3. $y = x - \tan x + \frac{1}{3} \tan^3 x$
4. $y = 3x + 2 \sin x \cos^3 x + 3 \sin x \cos x$

In Exercises 5–18, find the integral.

5. $\int \cos^5 x \sin x \, dx$ 6. $\int \cos^3 x \sin^4 x \, dx$

7. $\int \sin^7 2x \cos 2x \, dx$ 8. $\int \sin^3 x \, dx$

9. $\int \sin^3 x \cos^2 x \, dx$ 10. $\int \cos^3 \frac{x}{3} \, dx$

11. $\int \sin^3 2\theta \sqrt{\cos 2\theta} \, d\theta$ 12. $\int \frac{\cos^5 t}{\sqrt{\sin t}} \, dt$

13. $\int \cos^2 3x \, dx$ 14. $\int \sin^2 5x \, dx$

15. $\int \cos^4 3\alpha \, d\alpha$ 16. $\int \sin^4 6\theta \, d\theta$

17. $\int x \sin^2 x \, dx$ 18. $\int x^2 \sin^2 x \, dx$

In Exercises 19–24, use Wallis's Formulas to evaluate the integral.

19. $\int_0^{\pi/2} \cos^7 x \, dx$ 20. $\int_0^{\pi/2} \cos^9 x \, dx$

21. $\int_0^{\pi/2} \cos^{10} x \, dx$ 22. $\int_0^{\pi/2} \sin^5 x \, dx$

23. $\int_0^{\pi/2} \sin^6 x \, dx$ 24. $\int_0^{\pi/2} \sin^8 x \, dx$

In Exercises 25–42, find the integral involving secant and tangent.

25. $\int \sec 7x \, dx$ 26. $\int \sec^2(2x - 1) \, dx$

27. $\int \sec^4 5x \, dx$ 28. $\int \sec^6 3x \, dx$

29. $\int \sec^3 \pi x \, dx$ 30. $\int \tan^5 x \, dx$

31. $\int \tan^5 \frac{x}{2} \, dx$ 32. $\int \tan^3 \frac{\pi x}{2} \sec^2 \frac{\pi x}{2} \, dx$

33. $\int \sec^2 x \tan x \, dx$

35. $\int \tan^2 x \sec^4 x \, dx$

37. $\int \sec^6 4x \tan 4x \, dx$

39. $\int \sec^5 x \tan^3 x \, dx$

41. $\int \frac{\tan^2 x}{\sec x} \, dx$

34. $\int \tan^3 2t \sec^3 2t \, dt$

36. $\int \tan^5 2x \sec^4 2x \, dx$

38. $\int \sec^2 \frac{x}{2} \tan \frac{x}{2} \, dx$

40. $\int \tan^3 3x \, dx$

42. $\int \frac{\tan^2 x}{\sec^5 x} \, dx$


In Exercises 43–46, solve the differential equation.

43. $\frac{dr}{d\theta} = \sin^4 \pi\theta$

44. $\frac{ds}{d\alpha} = \sin^2 \frac{\alpha}{2} \cos^2 \frac{\alpha}{2}$



45. $y' = \tan^3 3x \sec 3x$

46. $y' = \sqrt{\tan x} \sec^4 x$

 **Slope Fields** In Exercises 47 and 48, a differential equation, a point, and a slope field are given. (a) Sketch two approximate solutions of the differential equation on the slope field, one of which passes through the given point. (b) Use integration to find the particular solution of the differential equation and use a graphing utility to graph the solution. Compare the result with the sketches in part (a). To print an enlarged copy of the graph, go to the website www.mathgraphs.com.

47. $\frac{dy}{dx} = \sin^2 x, (0, 0)$

48. $\frac{dy}{dx} = \sec^2 x \tan^2 x, (0, -\frac{1}{4})$



CAS **Slope Fields** In Exercises 49 and 50, use a computer algebra system to graph the slope field for the differential equation, and graph the solution through the specified initial condition.

49. $\frac{dy}{dx} = \frac{3 \sin x}{y}, y(0) = 2$

50. $\frac{dy}{dx} = 3\sqrt{y} \tan^2 x, y(0) = 3$

In Exercises 51–56, find the integral.

51. $\int \cos 2x \cos 6x \, dx$

52. $\int \cos 4\theta \cos(-3\theta) \, d\theta$

53. $\int \sin 2x \cos 4x \, dx$

54. $\int \sin(-4x) \cos 3x \, dx$

55. $\int \sin \theta \sin 3\theta \, d\theta$

56. $\int \sin 5x \sin 4x \, dx$

In Exercises 57–66, find the integral. Use a computer algebra system to confirm your result.

57. $\int \cot^3 2x \, dx$ 58. $\int \tan^4 \frac{x}{2} \sec^4 \frac{x}{2} \, dx$

59. $\int \csc^4 2x \, dx$ 60. $\int \cot^3 x \csc^3 x \, dx$

61. $\int \frac{\cot^2 t}{\csc t} \, dt$ 62. $\int \frac{\cot^3 t}{\csc t} \, dt$

63. $\int \frac{1}{\sec x \tan x} \, dx$ 64. $\int \frac{\sin^2 x - \cos^2 x}{\cos x} \, dx$

65. $\int (\tan^4 t - \sec^4 t) \, dt$ 66. $\int \frac{1 - \sec t}{\cos t - 1} \, dt$

In Exercises 67–74, evaluate the definite integral.

67. $\int_{-\pi}^{\pi} \sin^2 x \, dx$ 68. $\int_0^{\pi/3} \tan^2 x \, dx$

69. $\int_0^{\pi/4} 6 \tan^3 x \, dx$ 70. $\int_0^{\pi/4} \sec^2 t \sqrt{\tan t} \, dt$

71. $\int_0^{\pi/2} \frac{\cos t}{1 + \sin t} \, dt$ 72. $\int_{-\pi}^{\pi} \sin 5x \cos 3x \, dx$

73. $\int_{-\pi/2}^{\pi/2} 3 \cos^3 x \, dx$ 74. $\int_{-\pi/2}^{\pi/2} (\sin^2 x + 1) \, dx$

CAS In Exercises 75–80, use a computer algebra system to find the integral. Graph the antiderivatives for two different values of the constant of integration.

75. $\int \cos^4 \frac{x}{2} \, dx$ 76. $\int \sin^2 x \cos^2 x \, dx$

77. $\int \sec^5 \pi x \, dx$ 78. $\int \tan^3(1 - x) \, dx$

79. $\int \sec^5 \pi x \tan \pi x \, dx$ 80. $\int \sec^4(1 - x) \tan(1 - x) \, dx$

CAS In Exercises 81–84, use a computer algebra system to evaluate the definite integral.

81. $\int_0^{\pi/4} \sin 3\theta \sin 4\theta \, d\theta$ 82. $\int_0^{\pi/2} (1 - \cos \theta)^2 \, d\theta$

83. $\int_0^{\pi/2} \sin^4 x \, dx$ 84. $\int_0^{\pi/2} \sin^{12} x \, dx$

WRITING ABOUT CONCEPTS

85. In your own words, describe how you would integrate $\int \sin^m x \cos^n x \, dx$ for each condition.

- (a) m is positive and odd. (b) n is positive and odd.
- (c) m and n are both positive and even.

86. In your own words, describe how you would integrate $\int \sec^m x \tan^n x \, dx$ for each condition.

- (a) m is positive and even. (b) n is positive and odd.
- (c) n is positive and even, and there are no secant factors.
- (d) m is positive and odd, and there are no tangent factors.

87. Evaluate $\int \sin x \cos x \, dx$ using the given method. Explain how your answers differ for each method.

- (a) Substitution where $u = \sin x$
- (b) Substitution where $u = \cos x$
- (c) Integration by parts
- (d) Using the identity $\sin 2x = 2 \sin x \cos x$

CAPSTONE

88. For each pair of integrals, determine which one is more difficult to evaluate. Explain your reasoning.

- (a) $\int \sin^{372} x \cos x \, dx$, $\int \sin^4 x \cos^4 x \, dx$
- (b) $\int \tan^{400} x \sec^2 x \, dx$, $\int \tan^{400} x \sec x \, dx$



In Exercises 89 and 90, (a) find the indefinite integral in two different ways. (b) Use a graphing utility to graph the antiderivative (without the constant of integration) obtained by each method to show that the results differ only by a constant. (c) Verify analytically that the results differ only by a constant.

89. $\int \sec^4 3x \tan^3 3x \, dx$ 90. $\int \sec^2 x \tan x \, dx$

Area In Exercises 91–94, find the area of the region bounded by the graphs of the equations.

- 91. $y = \sin x$, $y = \sin^3 x$, $x = 0$, $x = \pi/2$
- 92. $y = \sin^2 \pi x$, $y = 0$, $x = 0$, $x = 1$
- 93. $y = \cos^2 x$, $y = \sin^2 x$, $x = -\pi/4$, $x = \pi/4$
- 94. $y = \cos^2 x$, $y = \sin x \cos x$, $x = -\pi/2$, $x = \pi/4$

Volume In Exercises 95 and 96, find the volume of the solid generated by revolving the region bounded by the graphs of the equations about the x -axis.

- 95. $y = \tan x$, $y = 0$, $x = -\pi/4$, $x = \pi/4$
- 96. $y = \cos \frac{x}{2}$, $y = \sin \frac{x}{2}$, $x = 0$, $x = \pi/2$

Volume and Centroid In Exercises 97 and 98, for the region bounded by the graphs of the equations, find (a) the volume of the solid formed by revolving the region about the x -axis and (b) the centroid of the region.

- 97. $y = \sin x$, $y = 0$, $x = 0$, $x = \pi$
- 98. $y = \cos x$, $y = 0$, $x = 0$, $x = \pi/2$

In Exercises 99–102, use integration by parts to verify the reduction formula.

99. $\int \sin^n x \, dx = -\frac{\sin^{n-1} x \cos x}{n} + \frac{n-1}{n} \int \sin^{n-2} x \, dx$

100. $\int \cos^n x \, dx = \frac{\cos^{n-1} x \sin x}{n} + \frac{n-1}{n} \int \cos^{n-2} x \, dx$

101. $\int \cos^m x \sin^n x \, dx = -\frac{\cos^{m+1} x \sin^{n-1} x}{m+n} + \frac{n-1}{m+n} \int \cos^m x \sin^{n-2} x \, dx$

102. $\int \sec^n x \, dx = \frac{1}{n-1} \sec^{n-2} x \tan x + \frac{n-2}{n-1} \int \sec^{n-2} x \, dx$

In Exercises 103–106, use the results of Exercises 99–102 to find the integral.

103. $\int \sin^5 x \, dx$

104. $\int \cos^4 x \, dx$

105. $\int \sec^4(2\pi x/5) \, dx$

106. $\int \sin^4 x \cos^2 x \, dx$

107. **Modeling Data** The table shows the normal maximum (high) and minimum (low) temperatures (in degrees Fahrenheit) in Erie, Pennsylvania for each month of the year. (Source: NOAA)

Month	Jan	Feb	Mar	Apr	May	Jun
Max	33.5	35.4	44.7	55.6	67.4	76.2
Min	20.3	20.9	28.2	37.9	48.7	58.5

Month	Jul	Aug	Sep	Oct	Nov	Dec
Max	80.4	79.0	72.0	61.0	49.3	38.6
Min	63.7	62.7	55.9	45.5	36.4	26.8

The maximum and minimum temperatures can be modeled by $f(t) = a_0 + a_1 \cos(\pi t/6) + b_1 \sin(\pi t/6)$, where $t = 0$ corresponds to January 1 and a_0, a_1 , and b_1 are as follows.


$a_0 = \frac{1}{12} \int_0^{12} f(t) \, dt$

$a_1 = \frac{1}{6} \int_0^{12} f(t) \cos \frac{\pi t}{6} \, dt$

$b_1 = \frac{1}{6} \int_0^{12} f(t) \sin \frac{\pi t}{6} \, dt$

(a) Approximate the model $H(t)$ for the maximum temperatures. (Hint: Use Simpson's Rule to approximate the integrals and use the January data twice.)

(b) Repeat part (a) for a model $L(t)$ for the minimum temperature data.

 (c) Use a graphing utility to graph each model. During what part of the year is the difference between the maximum and minimum temperatures greatest?

108. **Wallis's Formulas** Use the result of Exercise 100 to prove the following versions of Wallis's Formulas.

(a) If n is odd ($n \geq 3$), then

$\int_0^{\pi/2} \cos^n x \, dx = \left(\frac{2}{3}\right)\left(\frac{4}{5}\right)\left(\frac{6}{7}\right) \cdots \left(\frac{n-1}{n}\right).$

(b) If n is even ($n \geq 2$), then

$\int_0^{\pi/2} \cos^n x \, dx = \left(\frac{1}{2}\right)\left(\frac{3}{4}\right)\left(\frac{5}{6}\right) \cdots \left(\frac{n-1}{n}\right)\left(\frac{\pi}{2}\right).$

109. The **inner product** of two functions f and g on $[a, b]$ is given by $\langle f, g \rangle = \int_a^b f(x)g(x) \, dx$. Two distinct functions f and g are said to be **orthogonal** if $\langle f, g \rangle = 0$. Show that the following set of functions is orthogonal on $[-\pi, \pi]$.

$\{\sin x, \sin 2x, \sin 3x, \dots, \cos x, \cos 2x, \cos 3x, \dots\}$

110. **Fourier Series** The following sum is a *finite Fourier series*.

$f(x) = \sum_{i=1}^N a_i \sin ix$

$= a_1 \sin x + a_2 \sin 2x + a_3 \sin 3x + \cdots + a_N \sin Nx$


(a) Use Exercise 109 to show that the n th coefficient a_n is given by $a_n = (1/\pi) \int_{-\pi}^{\pi} f(x) \sin nx \, dx$.

(b) Let $f(x) = x$. Find a_1, a_2 , and a_3 .

SECTION PROJECT

Power Lines

Power lines are constructed by stringing wire between supports and adjusting the tension on each span. The wire hangs between supports in the shape of a catenary, as shown in the figure.




Let T be the tension (in pounds) on a span of wire, let u be the density (in pounds per foot), let $g \approx 32.2$ be the acceleration due to gravity (in feet per second per second), and let L be the distance (in feet) between the supports. Then the equation of the catenary is

$y = \frac{T}{ug} \left(\cosh \frac{ugx}{T} - 1 \right)$, where x and y are measured in feet.

(a) Find the length of the wire between two spans.

(b) To measure the tension in a span, power line workers use the *return wave method*. The wire is struck at one support, creating a wave in the line, and the time t (in seconds) it takes for the wave to make a round trip is measured. The velocity v (in feet per second) is given by $v = \sqrt{T/u}$. How long does it take the wave to make a round trip between supports?

(c) The sag s (in inches) can be obtained by evaluating y when $x = L/2$ in the equation for the catenary (and multiplying by 12). In practice, however, power line workers use the "lineman's equation" given by $s \approx 12.075t^2$. Use the fact that $[\cosh(ugL/2T) + 1] \approx 2$ to derive this equation.

 **FOR FURTHER INFORMATION** To learn more about the mathematics of power lines, see the article "Constructing Power Lines" by Thomas O'Neil in *The UMAP Journal*.